

**Grade 12 – APM Trials Exam 2015 - P2 memo**

1.1 A. Prove true for  $n = 1$

$$(1)^3 + 2(1) = 3 \text{ which is divisible by } 3. \\ \therefore \text{ True for } n = 1$$

B. Assume true for  $n = k$  where  $k \in \mathbb{N}$

$$k^3 + 2k \text{ is divisible by } 3$$

C. Prove true for  $n = k + 1$

$$(k + 1)^3 + 2(k + 1) \\ = k^3 + 3k^2 + 3k + 1 + 2k + 2 \\ = (k^3 + 2k) + 3k^2 + 3k + 3 \quad \text{since } k^3 + 2k \text{ is divisible by } 3$$

$$\therefore (k + 1)^3 + 2(k + 1) \text{ is divisible by } 3 \\ \therefore \text{ the statement is true for } n = k + 1$$

By the principle of Mathematical Induction  $n^3 + 2n$  is divisible by 3 for  $n \in \mathbb{N}$

(10)

1.2 
$$\frac{x - 10}{2x^2 + 5x - 3} = \frac{x - 10}{(2x - 1)(x + 3)}$$

$$\frac{x - 10}{(2x - 1)(x + 3)} = \frac{A}{2x - 1} + \frac{B}{x + 3}$$

$$\frac{x - 10}{(2x - 1)(x + 3)} = \frac{A(x + 3)}{2x - 1} + \frac{B(2x - 1)}{x + 3}$$

$$A + 2B = 1 \quad 3A - B = -10 \\ A = -\frac{19}{7} \quad B = \frac{13}{7}$$

$$\therefore \frac{x - 10}{(2x - 1)(x + 3)} = \frac{-19}{7(2x - 1)} + \frac{13}{7(x + 3)}$$

(8)

1.3  $x = 1 + i \quad \therefore x = 1 - i$

$$\text{sum} = 2 \quad \text{product} = 1 - i^2 = 2$$

$$\therefore x^2 - 2x + 2 \text{ is a factor of } x^3 + ax^2 + bx - 6$$

$$(x^2 - 2x + 2)(x - 3) = x^3 + 5x^2 + 8x + 6$$

$$\therefore a = -8 \quad \text{and } b = 5$$

(8)

$$\begin{aligned}
2.1 \quad & \frac{e^x}{e^x - 1} = 5 \\
& e^x = 5 \cdot e^x - 5 \\
& 4 \cdot e^x - 5 = 0 \\
& e^x = \frac{5}{4}
\end{aligned} \tag{4}$$


$$\begin{aligned}
2.2 \quad & \frac{-6}{1-x} \leq x \\
& \frac{-6}{1-x} - x \leq 0 \\
& \frac{-6 - x(1-x)}{1-x} \leq 0 \\
& \frac{x^2 - x - 6}{1-x} \leq 0 \\
& \frac{(x-3)(x+2)}{x-1} \geq 0 \\
& -2 \leq x < 1 \quad \text{or} \quad x > 3
\end{aligned} \tag{6}$$

$$\begin{aligned}
2.3 \quad & |x|^2 - 5|x| - 14 = 0 \\
& |x| = 7 \quad |x| \neq -2 \\
& \therefore x = \pm 7
\end{aligned} \tag{6}$$

$$\begin{aligned}
2.4 \quad & 3(\ln x)^2 + \ln x - 1 + \frac{1}{3(\ln x)^2 + \ln x - 3} = 0 \\
& \text{Let } k = 3(\ln x)^2 + \ln x \\
& k - 1 + \frac{1}{k-3} = 0 \\
& k^2 - 4k + 3 + 1 = 0 \\
& k^2 - 4k + 4 = 0 \\
& k = 2 \\
& 3(\ln x)^2 + \ln x - 2 = 0 \\
& (3 \ln x - 2)(\ln x + 1) = 0 \\
& \ln x = \frac{2}{3} \quad \ln x = -1 \\
& x = e^{\frac{2}{3}} \quad x = e^{-1}
\end{aligned} \tag{8}$$

$$\begin{aligned}
 3.1 \quad & \lim_{x \rightarrow \infty} \frac{x^2 - x + 6}{4x^3 + 3x^2 + x} \\
 & = 0
 \end{aligned}
 \tag{3}$$

$$\begin{aligned}
 3.2 \quad & \lim_{x \rightarrow \frac{\pi}{4}} \frac{\cos 2}{\cos \theta - \sin \theta} \\
 & = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\cos^2 \theta - \sin^2 \theta}{\cos \theta - \sin \theta} \\
 & = \lim_{x \rightarrow \frac{\pi}{4}} \frac{(\cos \theta + \sin \theta)(\cos \theta - \sin \theta)}{\cos \theta - \sin \theta} \\
 & = \lim_{x \rightarrow \frac{\pi}{4}} (\cos \theta + \sin \theta) \\
 & = \sqrt{2}
 \end{aligned}
 \tag{5}$$



4.1.2  $f(2) = 2 + 1 = 3$

$$\begin{aligned}
 \lim_{x \rightarrow 2^-} f(x) &= 2 + 1 = 3 \\
 \lim_{x \rightarrow 2^+} f(x) &= |2 - 4| = 2 \\
 \therefore \lim_{x \rightarrow 2^-} f(x) &\neq \lim_{x \rightarrow 2^+} f(x)
 \end{aligned}$$

Not continuous at  $x = 2$   
 Jump discontinuity.

(4)

4.2  $\lim_{x \rightarrow -1^-} f(x) = \lim_{x \rightarrow -1^+} f(x) = f(x)$

$$\begin{aligned}
 -a + b + 1 &= -1 - 3b \\
 a &= 4b + 2 \dots(1)
 \end{aligned}$$

$$\begin{aligned}
 ab &= -1 - 3b \\
 b(4b + 2) &= -1 - 3b \\
 4b^2 + 5b + 1 &= 0 \\
 b &= -\frac{1}{4} \quad \text{or} \quad b = -1 \\
 a &= 1 \quad \text{or} \quad a = -2
 \end{aligned}
 \tag{6}$$

$$5.1.1 \quad f \circ g(x) = \sqrt{(x^2 + 1)^2 - 1} \quad (4)$$

$$5.1.2 \quad \begin{aligned} (x^2 + 1)^2 - 1 &\geq 0 \\ x^4 + 2x^2 &\geq 0 \\ x^2(x^2 + 2) &\geq 0 \end{aligned}$$

$$x \in \mathbb{R} \quad (2)$$

$$5.1.3 \quad \begin{aligned} D_x[\sqrt{x+1} \cdot (x^2 + 1)^2] \\ = \frac{1}{2}(x+1)^{-\frac{1}{2}}(x^2 + 1)^2 + 2(x^2 + 1)(2x)(x+1)^{\frac{1}{2}} \end{aligned} \quad (6)$$

$$5.2.1 \quad f'(x) = 4(2x^2 + x - 1)^3(4x + 1) \quad (5)$$

$$5.2.2 \quad \begin{aligned} D_x \left[ \frac{\cos 3x}{\tan 5x} \right] \\ = \frac{(-3 \sin 3x \cdot \tan 5x - \sec 5x \cdot \tan 5x \cdot \cos 3x)}{\tan^2 x} \end{aligned} \quad (7)$$

$$6.1 \quad f(x) = (1 - x)^{-5}$$

$$f'(x) = -5(1 - x)^{-6}(-1)$$

$$f''(x) = (-5)(-6)(1 - x)^{-7}(-1)(-1)$$

$$f^n(x) = \frac{(n + 4)!}{4!} (1 - x)^{-(n+5)}$$

(6)

$$6.2 \quad xy^3 + 3x^2 = xy + 12$$

$$y^3 + 3xy^2 \frac{dy}{dx} + 6x = y + x \frac{dy}{dx}$$

$$3xy^2 \frac{dy}{dx} - x \frac{dy}{dx} = -3x - y^3 + y$$

$$\frac{dy}{dx}(3xy^2 - x) = -3x - y^3 + y$$

$$\frac{dy}{dx} = \frac{3x - y^3 + y}{3xy^2 - x} = -3$$

$$1 = -3(2) + c$$

$$y = -3x + 7$$

(8)

$$7.1 \quad x^2 - 2x - 3 = 0$$


$$x = -1 \quad x = 3$$

(4)

$$7.2 \quad y = \left(\frac{1}{3}x + \frac{2}{3}\right) \text{ by long division or other method.}$$

(6)

7.3



(8)

$$8.1 \quad 2 \cos \theta + \theta - 2 = 0$$

$$\theta = 3: 2 \cos 3 + 3 - 2 = -0,98$$

$$\theta = 5: 2 \cos 5 + 5 - 2 = 3,57$$

$\therefore$  Graph passes through the  $x$  - axis between 3 and 5 (4)

$$8.2 \quad f(x) = 2 \cos \theta + \theta - 2 = 0$$

$$f'(x) = -2 \sin \theta + 1$$

$$x_1 = 4$$

$$x_2 = 4 - \frac{f(4)}{f'(4)} = 3,72441$$

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)} = 3,69843$$

$$x_4 = x_3 - \frac{f(x_3)}{f'(x_3)} = 3,69815$$

$$x_5 = x_4 - \frac{f(x_4)}{f'(x_4)} = 3,69815$$

$\therefore x = 3,69815$  (6)

$$9.1 \quad \int x \sqrt{x^2 + 3} \, dx$$

$$= \int u^{\frac{1}{2}} \frac{du}{2}$$

$$\text{Let } u = x^2 + 3 \quad du = 2x \, dx$$

$$= \frac{1}{2} \times \frac{u^{\frac{3}{2}}}{\frac{3}{2}} + c$$

$$= \frac{(x^2 + 3)^{\frac{3}{2}}}{3} + c$$

(6)

$$9.2 \quad \int \sin 4x \cos 3x \, dx$$

$$= \frac{1}{2} \int (\sin 7x - \sin x) \, dx$$

$$= \frac{1}{2} \left( -\frac{\cos 7x}{7} - (-\cos x) + c \right)$$

$$= -\frac{\cos 7x}{14} + \cos x + c$$

(6)

$$9.3 \quad \int x \sin^2 x \, dx$$

$$= x \left( \frac{x}{2} - \frac{\sin 2x}{4} \right) - \int \left( \frac{x}{2} - \frac{\sin 2x}{4} \right) dx \quad \text{Let } u = x \quad dv = \sin^2 x \, dx = \frac{1}{2}(1 - \cos 2x) dx$$

$$= x \left( \frac{x}{2} - \frac{\sin 2x}{4} \right) - \left( \frac{x^2}{4} + \frac{\cos 2x}{8} \right) + c \quad du = 1 dx \quad v = \frac{x}{2} - \frac{\sin 2x}{4}$$

(9)

$$9.4 \quad \int \frac{x}{\sqrt{x-1}} \, dx$$

$$\text{Let } u = x - 1 \quad du = dx$$

$$= \int u^{-\frac{1}{2}}(u+1) du \quad x = u + 1$$

$$= \int \left( u^{\frac{1}{2}} + u^{-\frac{1}{2}} \right) du$$

$$= \frac{2(x-1)^{\frac{3}{2}}}{3} + 2(x-1)^{\frac{1}{2}} + c$$

(9)

$$10.1 \quad (x+2)^2 - 3 = 2x + 4$$

$$\begin{aligned} x &= -3 & x &= 1 \\ y &= -2 & y &= 6 \end{aligned}$$

(4)

$$10.2 \quad \text{Distance} = 2x + 4 - x^2 - 4x - 4 + 3 = -x^2 - 2x + 3$$

$$D_x[-x^2 - 2x + 3] = 0$$

$$-2x - 2 = 0$$

$$x = -1$$

$$\text{Max distance} = -(-1)^2 - 2(-1) + 3 = 4 \text{ units}$$

(6)

10.3

$$\text{Area} = \lim_{n \rightarrow \infty} \frac{4}{n} \sum_{i=1}^n f\left(\frac{4i}{n}\right)$$

$$= \lim_{n \rightarrow \infty} \frac{4}{n} \sum_{i=1}^n \left( -\left(\frac{4i}{n}\right)^2 - 2\left(\frac{4i}{n}\right) + 3 \right)$$

$$= \lim_{n \rightarrow \infty} \frac{4}{n} \sum_{i=1}^n \left( -\frac{16}{n^2} i^2 - \frac{8}{n} i + 3 \right)$$

$$= \lim_{n \rightarrow \infty} \frac{4}{n} \left( -\frac{16}{n^2} \left( \frac{n^3}{3} + \frac{n^2}{2} + \frac{n}{2} \right) - \frac{8}{n} \left( \frac{n^2}{2} + \frac{n}{2} \right) + 3n \right)$$

$$= \lim_{n \rightarrow \infty} \left( -\frac{64}{3} - \frac{32}{n} - \frac{32}{n^2} + 16 + \frac{16}{n} + 12 \right)$$

$$= \frac{88}{3}$$

(10)

$$\begin{aligned} 10.4 \quad & \pi \int_0^1 (-x^2 - 2x + 3)^2 dx \\ &= \pi \int_0^1 (x^4 + 4x^3 - 2x^2 - 12x + 9) dx \\ &= \pi \left( \frac{x^5}{5} + x^4 - \frac{2x^3}{3} - 6x^2 + 9x + c \right) \Big|_0^1 \\ &= \pi \left( \frac{1^5}{5} + (1)^4 - \frac{2(1)^3}{3} - 6(1)^2 + 9(1) \right) \\ &= \frac{53}{15} \pi \end{aligned}$$

(8)